Astronomy & Astrophysics manuscript no. aa3298
(DOI: will be inserted by hand later) February 2, 2008

arXiv:astro-ph/0506592v1 24 Jun 2005

On metal-deficient barium stars and their link with yellow symbiotic stars*

A. Jorissen[1**], L. Začs[2], S. Udry[3], H. Lindgren[4], and F.A. Musaev[5,6,7]

[1] Institut d'Astronomie et d'Astrophysique, Université Libre de Bruxelles, CP 226, Boulevard du Triomphe, B-1050 Bruxelles, Belgium
[2] Institute of Atomic Physics and Spectroscopy, University of Latvia, Raiņa bulvāris 19, Rīga, LV 1586 Latvia
[3] Observatoire de Genève, CH-1290 Sauverny, Suisse
[4] Lund Observatory, Box 43, SE-221 00, Lund, Sweden
[5] Special Astrophysical Observatory and Isaac Newton Institute of Chile, SAO Branch, Nizhnij Arkhyz, 369167, Russia
[6] ICAMER, National Academy of Sciences of Ukraine, 361605 Peak Terskol, Kabardino-Balkaria, Russia
[7] Shamakhy Astrophysical Observatory, National Academy of Sciences of Azerbaijan, Azerbaijan

Received / Accepted

Abstract. This paper addresses the question of why metal-deficient barium stars are not yellow symbiotic stars (YSyS). Samples of (suspected) metal-deficient barium (mdBa) stars and YSyS have been collected from the literature, and their properties reviewed. It appears in particular that the barium nature of the suspected mdBa stars needs to be ascertained by detailed abundance analyses. Abundances are therefore derived for two of them, HD 139409 and HD 148897, which reveal that HD 148897 should not be considered a barium star. HD 139409 is a mild barium star, with overabundances observed only for elements belonging to the first s-process peak (Y and Zr). It is only moderately metal-poor ([Fe/H] = −0.4). The evidence for binarity among metal-deficient barium stars is then reviewed, using three different methods: (i) radial-velocity variations (from CORAVEL observations), (ii) Hipparcos astrometric data, and (iii) a method based on the comparison between the Hipparcos and Tycho-2 proper motions. An orbit is obtained for HIP 55852, whereas evidence for the (so far unknown) binary nature of HIP 34795, HIP 76605, HIP 97874 and HIP 107478 is presented. No conclusion regarding the binary nature of HIP 11595, HIP 25161 could be reached. Two stars with no evidence for binarity whatsoever (HIP 58596 and BD +3°2688) are candidates low-metallicity thermally-pulsing asymptotic giant branch stars, as inferred from their large luminosities. The reason why mdBa stars are not YSyS is suggested to lie in their different orbital period distributions: mdBa stars have on average longer orbital periods than YSyS, and hence their companion accretes matter at a lower rate, for a given mass loss rate of the giant star. The definite validation of this explanation should nevertheless await the determination of the orbital periods for the many mdBa stars still lacking periods, in order to make the comparison more significant.

Key words. binaries: symbiotic - Stars: abundances - Stars: AGB and post-AGB - binaries: spectroscopic

1. The problem

Our understanding of the link between chemically-peculiar red giants like barium stars or CH stars, and yellow symbiotic stars (YSyS) has made substantial progress in the last decade (see the reviews by Jorissen, 2003a,b), mainly with the realisation that likely all yellow symbiotics (i.e., involving a giant of spectral type G or K as primary component) involve barium stars (Smith et al., 1996, 1997; Pereira & Porto de Mello, 1997; Pereira et al., 1998). The metal-deficient nature of the giant is a key factor, because it implies a rather large luminosity for the giant. Since evolutionary tracks of metal-deficient stars are shifted towards the blue, metal-deficient giants of spectral type K must lie on the upper part of the (asymptotic) giant branch (see Fig. 11 of Smith et al., 1996), where they suffer strong mass loss. If such metal-deficient K giants are in binary systems, their strong wind will interact with the companion and trigger symbiotic activity.

The other facet of this problem, namely whether all metal-deficient barium stars are symbiotic stars, is not

---

[*] Based on observations carried out at the European Southern Observatory (ESO, La Silla, Chile) and with the 1-m Swiss telescope at the Haute-Provence Observatory

[**] Senior Research Associate, F.N.R.S., Belgium

yet fully answered. The present paper offers a first step in that direction.

We have collected a list of candidate metal-deficient barium stars and have assembled new observations to check (i) whether these stars are binaries, (ii) whether they are barium stars and (iii) whether they exhibit symbiotic activity.

2. The samples

Before discussing metal-deficient barium stars, it is useful to first summarize the properties of YSyS, to which metal-deficient barium stars may be compared.

2.1. Yellow symbiotic stars

All known YSyS are listed in Table 1, which shows that all the stars studied so far exhibit the barium syndrome. YSyS with a *stellar* infrared continuum (s-type, as opposed to the dusty d'-type; see below) are clearly metal-deficient objects, as revealed by their low metallicities and high space velocities (CD −43°14304 may be an exception; however, it is of spectral type K7, and should perhaps not be included in the family of YSyS). The presence of the barium syndrome among a family of binary metal-deficient stars fully supports the commonly accepted hypothesis that the s-process is more efficient at low metallicities (Clayton, 1988; Jorissen, 2003a). s-Type YSyS, with their metallicities lower than classical barium stars, may be expected to be, on average, more luminous than the latter (see Fig. 11 of Smith et al., 1996, comparing the luminosity function of Pop.I and Pop.II K giants). This is a direct consequence of the fact that evolutionary tracks shift towards the blue in the Hertzsprung-Russell (HR) diagram as metallicity decreases, as shown in Fig. 1b. Fig. 1a confirms that the YSyS AG Dra and BD −21°3873 are indeed more luminous than classical barium stars. This difference in the average luminosity – and hence mass-loss rate – of the two populations thus explains why YSyS, despite hosting a K giant, exhibit symbiotic activity whereas barium stars do not. The larger mass-loss rates for the cool components of s-type YSyS – as compared to Ba stars – may be inferred from the comparison of their IRAS [12] − [25] color indices, which reflect the amount of dust present in the system: $([12] - [25])_{\rm Ba} < 0.1$, as compared to 0.45 for AG Dra (Smith et al., 1996). Mürset et al. (1991) and Drake et al. (1987) provide direct measurements (or upper limits) for the mass loss rates of AG Dra and of Ba stars, respectively, which confirm the above conclusion.

YSyS with a *dusty* infrared continuum (d'-type; Allen, 1982; Schmid & Nussbaumer, 1993) differ from their s-type counterparts in several respects (Table 1): they host a complex circumstellar environment (including cool dust, bipolar outflows, extended optical nebulae or emission-line spectra closely resembling those of planetary nebulae), the cool components have early spectral types (F to early K), they are often fast rotators (with the possible exception of M 1-2 =V471 Per; Grauer & Bond, 1981) and, finally, they belong to the galactic disk unlike s-type YSyS which belong to the halo.

All these arguments suggest that the hot component in d'-type SyS has just evolved from the AGB to the WD stage. The rather cool dust (Schmid & Nussbaumer, 1993) is a relic from the mass lost by the AGB star. The optical nebulae observed in d'-type SyS are most likely genuine planetary nebulae rather than the nebulae associated with the ionized wind of the cool component (Corradi et al., 1999). This is especially clear for AS 201 which actually hosts *two* nebulae (Schwarz, 1991): a large fossil planetary nebula detected by direct imaging, and a small nebula formed in the wind of the current cool component. Finally, the rapid rotation of the cool component has likely been caused by spin accretion from the former AGB wind like in WIRRING systems (Jeffries & Stevens, 1996; Jorissen, 2003b). The fact that the cool star has not yet been slowed down by magnetic braking is another indication that the mass transfer occurred fairly recently (Theuns et al., 1996). Corradi & Schwarz (1997) obtained 4000 y for the age of the nebula around AS 201, and 40000 y for V417 Cen.

2.2. Metal-deficient barium stars

Metal-deficient barium stars (with metallicities in the range −1.1 to −1.8, comparable to that of YSyS) were identified by Luck & Bond (1991), Mennessier et al. (1997) and Začs et al. (2000), and occupy the same region of the HR diagram as YSyS (Fig. 1b). The question thus arises why metal-deficient barium stars are not SyS. Different answers must be sought, depending upon their absolute visual magnitudes $M_{\rm V}$. The most luminous systems, with $M_{\rm V} < -2$, are likely located on the thermally-pulsing AGB, so that their Ba syndrome may be explained by internal nucleosynthesis. They thus should not be binaries, and therefore cannot be SyS! HD 104340 (open circle in Fig. 1b), a metal-deficient Ba star studied by Junqueira & Pereira (2001), and BD +03°2688 (Table 2) provide good illustrations of this situation, since they both lie above the TP-AGB threshold and CORAVEL radial-velocity measurements spanning several years do not reveal any clear orbital motion (Figs. 9 and 10, as well as Sect. 4).

The less luminous and warmest among metal-deficient Ba stars, clumping around $M_{\rm V} \sim +1$ in the HR diagram, are also sometimes classified as CH stars (crosses in Fig. 1b). They are not losing mass at a large enough rate to trigger any symbiotic activity, as revealed by their small [12] − [25] color indices (< 0.3; Smith et al., 1996).

Finally, at intermediate luminosities ($-2 \le M_{\rm V} \le +1$) where YSyS are located, metal-deficient Ba stars are not luminous enough to be TP-AGB (hence they should be binaries), but yet their mass loss rates must be large enough to trigger symbiotic activity (provided that the orbital separation is not too large, since it is the mass *accretion* rate by the compact companion which is in fact the key pa-

Table 1. The barium syndrome among YSyS. The spectral type of the cool component is taken from Mürset & Schmid (1999), or references therein. In column labelled 'nebula', 'y' means that an optical nebula has been detected, and 'PN' that, based on its emission line spectrum, the star has traditionally been included in planetary nebulae catalogues, even though no optical nebula may be visible. The column labelled 'P' lists the orbital period, from Mürset & Schmid (1999)

Name	Sp. Typ.	[Fe/H]	V_r (km/s)	b (°)	[Ba/Fe]	$V \sin i$ (km/s)	nebula	P (d)	Ref.
					d'-type				
V417 Cen	G8-K2	~ 0.0		-1	0.5	70	y	247	(5,11)
HDE 330036 =Cn 1-1	G5	0.02	-14	$+4$	0.88	100	PN	-	(14,17)
AS 201	G5	0.07		$+7$	0.63	25	y	-	(12,17)
V471/V741 Per =M 1-2	G5	?	-12	-9	> 0		PN	-	(2)
St H$_\alpha$ 190	G5	0.0	~ 10	-35	~ 0.5	100	bip. outf.	-	(10,13)
Wray 157	G5	?							
Hen 1591	$<$ K4	?							
					s-type				
UKS Ce-1	C4,5Jch	?	$+20$	$+20$	> 0			-	(6)
S 32	C1,1CH	?	$+325$	-30	> 0			612	(6,14)
Hen 2-467	K0	-1.1	-109	-12	+0.8		n	478	(4,16)
BD$-21°$3873	K2	-1.1	$+204$	$+37$	+0.5		n	282	(3,15,16)
		-1.3			+0.3				(9)
AG Dra	K2	-1.3	-148	$+41$	+0.5		n	554	(8,16)
CD $-43°$14304	K7	-1.4	$+27$	-41	?			1448	(7,18)
					Chemical evolution of the Galaxy				
		-1.0			< 0.2				(1)

References: (1) Edvardsson et al., 1993, A&A, 275, 101 (2) Grauer & Bond, 1981, PASP, 93, 630 (3) Pereira et al., 1997, AJ, 114, 2128 (4) Pereira et al., 1998, AJ, 116, 1977 (5) Pereira et al., 2003, In: Symbiotic stars probing stellar evolution, R.L.M. Corradi, J. Mikołajewska, T.J. Mahoney (eds.), Astron. Soc. Pacific Conf. Ser. (San Francisco), p. 85 (6) Schmid, 1994, A&A, 284, 156 (7) Schmid et al., 1998, A&A, 329, 986 (8) Smith et al., 1996, A&A, 315, 179 (9) Smith et al., 1997, A&A, 324, 97 (10) Smith et al., 2001, ApJ, 556, L55 (11) Van Winckel et al., 1994, A&A, 285, 241 (12) Schwarz, 1991, A&A, 243, 469 (13) Munari et al., 2001, A&A, 369, L1 (14) Schmid & Nussbaumer, 1993, A&A, 268, 159 (15) Munari & Patat, 1993, A&A, 277, 195 (16) Corradi et al., 1999, A&A, 343, 841 (17) Pereira et al., 2005, A&A 429, 993 (18) The metallicity is from Pereira, priv. comm.

rameter; see Sect. 5 and Jorissen 2003a). It is thus of great interest to check (i) the Ba nature of those metal-deficient stars with intermediate luminosities, (ii) their binary nature, and (iii) their suspected symbiotic activity. The first two issues are addressed in Sects. 3 and 4, respectively.

As far as a possible symbiotic activity is concerned, there is no indication from their photometric $U-B$ and $B-V$ indices that the metal-deficient stars in Table 2 have a strong blue continuum which could betray their symbiotic nature. It is thus very likely that none among these stars is a full-fledged symbiotic star. No signature of weak symbiotic activity (of the kind exhibited by some binary S stars; see Fig. 14 and Van Eck & Jorissen, 2002) is observed in the H$_\alpha$ line profile either (Fig. 2).

3. Abundances

The classification of the stars in Table 2 as metal-deficient Ba stars is subject to caution, as it does not rely on spectroscopic abundance analyses, but rather on a maximum-likelihood assignment based on kinematic, spatial and luminosity properties (Mennessier et al., 1997) for barium stars from the list of Lü (1991). Nevertheless, when a metallicity determination is available, it confirms



Fig. 1. Left panel (a): Evolutionary tracks of Schaller et al. (1992) compared with the locations of classical barium stars (stars labelled ‘G’ in Mennessier et al. (1997); open dots) and the yellow SyS AG Dra (filled circle), BD −21°3873 (filled triangle) and Hen 2-467 (filled square). The bolometric magnitudes were taken from the references listed in Table 1. These bolometric magnitudes were combined with bolometric corrections from Bessell et al. (1998) and $B-V$ indices from Munari et al. (1992) and Munari & Buson (1992) to yield the absolute visual magnitudes.
Right panel (b): Same as (a) but for YSyS (filled symbols as in the left panel) and metal-deficient barium stars [open triangles: stars flagged as ‘H’ by Mennessier et al. (1997); crosses: CH stars also flagged as ‘H’ by Mennessier et al. (1997); small open squares: additional metal-deficient, s-process-rich stars from Table 2; open circle: HD 104340, large open square: HD 206983 from Junqueira & Pereira (2001)]. The dashed horizontal line represents the luminosity ($M_{\rm bol} = -3$, corresponding to $M_{\rm V} \sim -2$) at the first thermal pulse in a 1 M$_\odot$ AGB star of metallicity [Fe/H] =−1.8 according to Lattanzio (1991).

the metal-deficient nature of the object (see Table 2). HD 139409 is an exception, though, since detailed spectral analyses reveal that it is neither metal-poor nor strongly enriched in s-process elements (see below). It may nevertheless be hoped that the metal-deficient assignment made by Mennessier et al. (1997) is valid in all the other cases. Regarding the Ba nature of these stars, it is known that the Lü (1991) catalogue of barium stars, from which the sample of barium stars used by Mennessier et al. (1997) was drawn, is contaminated by many non-barium stars (Griffin & Keenan, 1992; Jorissen et al., 1996), especially among those stars having a Ba index smaller than 1.

It would therefore in principle be necessary to re-evaluate the Ba nature of all the stars listed in Table 2. So far, spectra could be obtained for two of them, HD 139409 and HD 148897, which are discussed in detail in the present section. Of these, only HD 139409 appears to be a mild barium star, thus confirming the suspicion about the Lü (1991) catalogue expressed above.

3.1. The case of HD 148897 and HD 139409

HD 148897 (= HR 6152) has been tagged as a ‘likely marginal barium star’ by Boyle & McClure (1975) and as a marginal CH star by Vilhu et al. (1977). It therefore found its way into the barium-star catalogs of Lü et al. (1983) and Lü (1991), as well as the catalog of CH and metal-deficient Ba stars of Šleivyte & Bartkevičius (1990). The star was classified as G8.5III CN-2 Fe-1 CH-1 by Keenan & McNeil (1989), and this classification as CN- and CH-weak contradicts the earlier assignments.

A detailed abundance analysis has thus been performed to clarify the situation, and its results are compared with previous analyses by Kyröläinen et al. (1986) and Luck (1991) in Sect. 3.1.2.

HD 139409 (= HIP 76605) has been classified as a marginal barium star by MacConnell et al. (1972), as G5 III Ba1 by Yamashita & Norimoto (1981) and as K0 III/II Ba 0.5 by Lü (1991).

3.1.1. Observations

A high-resolution spectrum of HD 148897 was obtained using the Coudé Matrix Echelle Spectrometer

Table 2. Stars classified as metal-deficient barium stars by Mennessier et al. (1997). The absolute magnitude M_V is a maximum likelihood estimate obtained by Mennessier et al. (1997), except for the additional stars where it is derived from a straight inversion of the Hipparcos parallax (BD +75°348), from a spectroscopic estimate of gravity and an educated guess for the mass (BD+3°2688) or from a fit to the M 92 isochrone (CS stars). The column labelled 'Ba' indicates whether detailed chemical analyses have confirmed the Ba nature of the star.

HIP	HD/DM	M_V	$B-V$	[Fe/H]	Ref	Ba	Ref	Rem
4347	5424	-0.6	1.14					
11595	15589	-0.5	1.15	-0.7	9			
25161	$-27°2233$	-1.1	1.13					
29740	43389	-3.5	1.48					
34795	55496	-1.7	0.96	-1.55	5	y	5	
58596	104340	-1.9	1.36	-1.72	3	y	3	
69834	123396	-0.9	1.19					
76605	139409	-0.5	1.04	-0.42	2	mild	2	$M_V = 1.5$ is derived from spectroscopic $\log g$
80843	148897	-1.8	1.27	-1.0	2	**n**	2	
				-0.62	7			
				-1.16	8			
97874	187762	-1.2	1.10					
107478	206983	0.2	1.64	-1.43	3	y	3	

Additional stars [from refs. (1), (5) and (10)]

HIP	HD/DM	M_V	$B-V$	[Fe/H]	Ref	Ba	Ref	Rem
43042	+75°348	-0.4 :	1.17	-0.8	1	y	1	
55852	+4°2466	?	0.60	-1.85	5,11	y	4,5	
	+3°2688	-5.0	1.22	-1.42	5,11	y	5	
	CS 22942-019	0	0.74	-2.67	10	y	10	
	CS 22948-027	-1	0.93	-2.60	10	y	10	

References to the Table:
(1) Začs et al. (2000) (2) This work (3) Junqueira & Pereira (2001) (4) Burris et al. (2000) (5) Luck & Bond (1991) (6) Jorissen et al. (1998) (7) Kyröläinen et al. (1986) (8) Luck (1991) (9) Barbuy et al. (1992) (10) Preston & Sneden (2001) (11) $B-V$ from the Tycho-2 catalog (Høg et al., 2000b)

(MAESTRO; Musaev et al., 1999) delivering a resolving power of 45 000 and installed on the 2-m Zeiss telescope of the Terskol Observatory (located in Northern Caucasus at an altitude of 3100 m). The spectrometer is equipped with a Wright Instruments CCD detector with 1242 × 1152 pixels (22.5 × 22.5 μm). A total exposure of 1800 s was taken on February 18, 2003. The spectrum covers the range 365 to 1020 nm spread over 85 spectral orders.

A high-resolution spectrum of HD 139409 was obtained on the HARPS spectrograph (Mayor et al., 2003), delivering a resolution of 115 000 and installed on the ESO La Silla 3.6 m telescope. A total exposure of 200 s was obtained by Xavier Bonfils on May 31, 2004. In order to check equivalent widths delivered by HARPS, a spectrum of the standard star Arcturus was obtained as well by Fabien Carrier. The HARPS spectra were reduced by the observers using standard pipeline processing. Equivalent widths for the same set of lines as those studied in HD 139409 have been measured by one of us (L.Z.) in the HARPS spectrum of Arcturus and compared to those from the Arcturus spectral atlas (Hinkle et al., 2000). The agreement between the two sets of equivalent widths is excellent (Fig. 3), thus qualifying HARPS for abundance analyses. Spectra around the λ614.172 nm Ba II line are shown in Fig. 4 for the two analyzed stars and for Arcturus (Hinkle et al., 2000).

3.1.2. Reduction and analysis

HD 148897

The reduction of the CCD frames (subtraction of bias, dark and scattered light, flat fielding, extraction of echelle orders and wavelength calibration) was performed with the DECH20T software (Galazutdinov, 1992). More than 500 weak to medium-strong atomic lines, free of blends, were identified and their equivalent widths were measured with the DECH routines. The radial velocity was measured using a large number of symmetric absorption lines. The observed velocity has been brought to the heliocentric system by adding +22.7 km s^{-1}. The mean heliocentric radial velocity for HD 144897 was found to be +16.7 km s^{-1}. The atmospheric parameters for that star cannot be derived from photometry, since the standard temperature calibrations refer to stars of normal chemical composition. To obtain a colour-independent estimate of the temperature, a spectroscopic temperature has been derived from the excitation equilibrium of Fe I lines. The surface gravity $\log g$ was determined using the FeI/FeII ionization balance, whereas the microturbulent velocity v_t was derived



Fig. 2. H_α profiles for two candidate metal-deficient Ba stars (HD 139409 and HD 148897; see however the discussion in Sect. 3), compared to that of the K giant Arcturus. The wavelength scale has been corrected for the stellar radial velocity. The profiles show no indication of symbiotic activity.

by forcing the abundances of individual Fe I lines to be independent of the reduced equivalent width. The resulting atmospheric parameters for HD 148897 are as follows: $T_{\rm eff}$ = 4350 K, $\log g = 1.0$ (cgs), and $v_t = 2.0$ km s^{-1}. An independent determination of the surface gravity, using the above $T_{\rm eff}$ value, Mennessier et al. (1997) absolute visual magnitude ($M_V = -1.8$), a bolometric correction of -0.5 and a mass of 1 M$_\odot$, yields $\log g = 1.1$, in agreement with the adopted value (see Luck, 1991, for a discussion of the discrepancy usually observed between the photometric and spectroscopic gravities). Comparison with atmospheric parameters from the literature is presented in Table 3.

The abundance analysis has been performed with the standard LTE line analysis program WIDTH9 developed by Kurucz. The model atmospheres were taken from Gustafsson et al. (1975). The synthetic spectra were generated using the spectral synthesis code STARSP (Tsymbal, 1996). Oscillator strengths have been taken from the VALD database (Piskunov et al., 1995). The resulting abundances (normalized by the solar-system abundances of Grevesse & Sauval, 1998) are listed in Table 3,



Fig. 3. Comparison of equivalent widths measured on the HARPS spectrum of Arcturus with those of the Arcturus spectral atlas (Hinkle et al., 2000), for the same set of lines as that measured in HD 139409.

from which it may be concluded that HD 148897 appears to be a rather typical metal-deficient star (McWilliam, 1997), and should certainly *not be considered as a (metal-deficient) barium star*.

Interestingly enough, there are no indications whatsoever from the Hipparcos and Tycho data (applying the methods described in Sect. 4) that this star is binary, in agreement with the fact that it is not a barium star.

HD 139409

The effective temperature of HD 139409 has been derived from the excitation equilibrium of Fe I, Ti I and Cr I lines (see Fig. 5 for Fe). The surface gravity $\log g$ was determined from the Fe I/Fe II ionization balance, and the microturbulent velocity by forcing the abundances of individual Fe I, Ti I and Cr I lines to be independent of equivalent width (see Fig. 6 for Fe). Although spectroscopic gravity and temperature determinations in late-type, metal-deficient stars are probably affected by non-LTE effects, these effects remain small when [Fe/H] ≥ -1.0 (see, for example, Allende Prieto et al., 1999). The stellar parameters of Arcturus ([Fe/H]$= -0.6$, $\log g = 1.3$) derived by the spectroscopic method (applied on the HARPS spectrum) are in good agreement indeed with those derived by other (non-spectroscopic) methods.

The resulting atmospheric parameters for HD 139409 are as follows: $T_{\rm eff}$ = 5000 K, $\log g = 2.8$ (cgs), and ξ_t = 2.0 km s^{-1}. The spectroscopic gravity, combined with a mass of 1 M$_\odot$, leads to $M_V \simeq +1.5$. Thus our calculations indicate that HD 139409 is less luminous than predicted

Table 3. Averaged absolute abundances (in the scale where $\log\epsilon(\mathrm{H}) = 12$) and normalized abundances [X/Fe] for HD 148897 (relative to the Sun; Grevesse & Sauval, 1998). The standard deviations σ and the number of lines used in the analysis (n) are also given. Two sets of abundances from Luck (1991) are given: one is based on a 'photometric $\log g$' (derived from the absolute magnitude, the effective temperature and an estimate for the stellar mass) and the other on the 'spectroscopic $\log g$' (Fe I/Fe II ionization balance).

	This work					Kiröläinen et al. (1986)		Luck (1991)		Luck (1991)	
	T_{eff}= 4350 K $\log g = 1.0$ $v_t = 2.0$ km s^{-1}					T_{eff}= 4360 K $\log g = 1.5$ $v_t = 2.6$ km s^{-1}		T_{eff}= 4100 K phot. $\log g = 1.6$ $v_t = 1.7$ km s^{-1}		T_{eff}= 4100 K spec. $\log g = 0.1$ $v_t = 1.7$ km s^{-1}	
X	$\log\epsilon$(X)	σ	n	$[\mathrm{X/H}]_\odot$	$\left[\frac{\mathrm{X}_{I,II}}{\mathrm{Fe}_{I,II}}\right]_\odot$	$[\mathrm{X/H}]_\odot$	$[\mathrm{X/Fe}]_\odot$	$[\mathrm{X/H}]_\odot$	$\left[\frac{\mathrm{X}_{I,II}}{\mathrm{Fe}_{I,II}}\right]_\odot$	$[\mathrm{X/H}]_\odot$	$\left[\frac{\mathrm{X}_{I,II}}{\mathrm{Fe}_{I,II}}\right]_\odot$
O I	8.22	0.20	4	-0.61	+0.41	-0.58	+0.04	-0.50	+0.50	-1.10	+0.06
Na I	5.10	0.08	6	-1.23	-0.21	-1.11	-0.49	-1.36	-0.36	-1.26	-0.10
Mg I	6.92	0.09	6	-0.66	+0.36	-0.59	+0.03	-0.54	+0.46	-0.62	+0.54
Al I	5.61	0.23	4	-0.86	+0.16	-0.70	-0.08	-1.28	-0.28	-1.20	-0.04
Si I	6.91	0.11	23	-0.77	+0.25	-0.56	+0.06	-0.24	+0.76	-0.59	+0.57
K I	4.17		1	-0.95	+0.07	-	-	-	-	-	-
Ca I	5.65	0.11	21	-0.71	+0.31	-0.54	+0.08	-1.09	-0.09	-0.97	+0.19
Ca II	5.48		1	-0.88	+0.09	-	-	-	-	-	-
Sc II	2.08	0.12	13	-1.09	-0.12	-0.86	-0.24	-0.75	-0.40	-1.37	-0.21
Ti I	4.20	0.07	56	-0.82	+0.20	-0.48	+0.14	-1.20	-0.20	-1.19	-0.03
Ti II	4.22	0.14	8	-0.80	+0.17	-	-	-0.51	-0.16	-1.16	0.00
V I	3.11	0.09	23	-0.89	+0.13	-0.56	+0.06	-1.24	-0.24	-1.24	-0.08
V II	3.01		1	-0.99	-0.02	-	-	-	-	-	-
Cr I	4.59	0.12	27	-1.08	-0.06	-0.82	-0.20	-1.19	-0.19	-1.08	+0.08
Cr II	4.85	0.08	6	-0.82	+0.15	-	-	-	-	-	-
Mn I	4.24	0.10	11	-1.15	-0.13	-1.00	-0.38	-	-	-	-
Fe I	6.48	0.07	180	-1.02	-	-0.62	-	-1.00	-	-1.16	-
Fe II	6.53	0.07	17	-0.97	-	-	-	-0.35	-	-1.16	-
Co I	4.11	0.12	21	-0.81	+0.21	-0.70	-0.08	-0.84	+0.16	-1.10	+0.06
Ni I	5.15	0.11	54	-1.10	-0.08	-0.62	0.00	-0.84	+0.16	-1.12	+0.04
Cu I	3.31	0.17	3	-0.90	+0.12	-	-	-0.64	+0.36	-0.80	+0.36
Zn I	3.78		1	-0.82	+0.20	-	-	-0.39	+0.61	-0.92	+0.24
Y II	1.21	0.12	6	-1.03	-0.06	-0.73	-0.11	-0.62	-0.27	-1.14	+0.02
Zr I								-1.43	-0.43	-1.43	-0.27
Zr II	1.91	0.13	3	-0.69	+0.28	-0.37	+0.25	-	-	-	-
Ba II	0.97	0.05	3	-1.16	-0.19	-0.66	-0.04	-	-	-	-
La II	0.24	0.25	3	-0.93	+0.04	-0.52	+0.10	-	-	-	-
Ce II	0.26	0.16	5	-1.32	-0.35	-0.65	-0.03	-0.68	-0.33	-1.21	-0.05
Pr II	-0.20	0.13	2	-0.91	+0.06	-	-	-	-	-	-
Nd II	0.56	0.13	8	-0.94	+0.03	-	-	-0.61	-0.26	-1.15	+0.01
Eu II	0.17		1	-0.34	+0.63	-	-	-0.40	-0.05	-1.01	+0.15

by Mennessier et al. (1997). The derived iron abundance is $[\mathrm{Fe/H}] = -0.42$ [adopting $\log(\epsilon(\mathrm{Fe})) = 7.50$].

The basic conclusion from the set of abundances listed in Table 4 and displayed in Fig. 7 is that HD 139409 appears to be a mild barium star.[1] The s-process overabundances observed in HD 139409, although quite moderate, are not much smaller than those observed in a yellow symbiotic star like BD -21°3873, which exhibit overabundances of s-process elements in the range 0.3 – 0.8 dex (Smith et al., 1997). With $[\mathrm{Fe/H}] = -0.4$, HD 139409 is, however, not as metal-deficient as the other stars considered in this paper.

4. Statistics of binarity among metal-deficient Ba stars

The list of confirmed or suspected metal-deficient barium stars remaining after the screening process based on abundances (as described in Sect. 3) is given in Table 5. The possible binary nature of those stars may be assessed using three different methods:

- checking for radial-velocity variations;
- checking for astrometric orbital motion, directly from the Hipparcos astrometric data;

---

[1] Interestingly, Pinsonneault et al. (1984) quoting a private communication from E. Luck, remark that Luck "is completing an abundance analysis of HD 139409, which will show that the light s-process elements do have ($\sim +1.5$ dex) enhancements, while heavier s-process element enhancements are much smaller ($\sim +0.3$ dex)". Luck's study seems to have never been published.



Fig. 4. The spectra of the target stars in the region of the Ba II line at 614.172 nm. Measured lines are marked. Also shown is the spectrum of the standard star Arcturus (Hinkle et al., 2000).



Fig. 5. The iron abundances derived from Fe I (open circles) and Fe II (filled circles) lines are displayed as a function of the excitation potential for the lower energy level of the line. The absence of a trend in these data is used to derive the spectroscopic temperature of HD 139409.



Fig. 6. The abundances derived from the Fe I (open circles) and Fe II (filled circles) lines are displayed as a function of the line equivalent widths. The absence of a trend in these data is used to derive the microturbulent velocity of HD 139409.



Fig. 7. Abundance pattern in HD 139409 (abundances derived from ionized species are represented by black dots, and from neutral species by crosses).

– checking for astrometric orbital motion, indirectly from a comparison of the Hipparcos and Tycho-2 proper motions.

These methods are now described in turn.

Table 5. Binary properties of confirmed or suspected metal-deficient barium stars. The column labeled 'Ba' indicates whether detailed chemical analyses have confirmed the Ba nature of the star. The columns labeled $\chi^2_{\mu_{\rm HIP}-\mu_{\rm Tyc}}$ and Prob($\chi^2 > \chi^2_{\mu_{\rm HIP}-\mu_{\rm Tyc}}$) provide the χ^2 (and its associated probability) involved in the comparison of Hipparcos and Tycho-2 proper motions (see text for details). The column 'Bin.' has been set to 'y' if the first kind risk of rejecting the null hypothesis that the Tycho and Hipparcos proper motions are equal is smaller than 10%. The column labeled IAD indicates whether the signature of an orbital motion is present in the Hipparcos Intermediate Astrometric Data, according to the various tests described in the text. The column labeled $\sigma(Vr)$ provides the radial-velocity standard deviation, Δt and N correspond to the time span and number of observations, respectively. The columns labeled P and 'Ref' list the orbital period (when available) and the reference for the radial velocity and/or orbital data. The column 'Binary' gives the final binarity diagnostics.

HIP	HD/DM	Ba	Ref	ϖ (mas)	$\mu_{\rm HIP}-\mu_{\rm Tyc}$			IAD	$\sigma(Vr)$	Δt	N	P	Ref	Binary
					$\chi^2_{\rm obs}$	Prob($\chi^2 > \chi^2_{\rm obs}$)	Bin.		(km s^{-1})	(d)		(d)		
4347	5424			0.22 ± 1.42	1.05	0.59	n	n	2.32	3306	13	1881	6	y
11595	15589			2.03 ± 1.21	1.63	0.44	n	n	-	-	-	-		?
25161	−27°2233			0.89 ± 1.35	0.23	0.89	n	n	-	-	-	-		?
29740	43389			−1.25 ± 1.00	4.54	0.10	y	y	3.85	3350	24	1689	6	y
34795	55496	y	5	2.44 ± 1.04	0.02	0.99	n	y	0.73	5121	24	-	2	y?
58596	104340	y	3	−0.97 ± 1.09	3.39	0.18	n	n	1.64	2587	16	-	2	n
69834	123396			1.73 ± 0.86	1.98	0.27	n	n	-	-	-	-		?
76605	139409	y	2	5.51 ± 1.14	4.98	0.08	y	n	0.66	1478	2	-	2	y?
97874	187762			2.07 ± 1.53	9.00	0.01	y	y	-	-	-	-		y
107478	206983	y	3	3.75 ± 1.86	2.94	0.23	n	y	-	-	-	-		y?

Additional stars [from refs. (1), (5) and (10)]

43042	+75°348	y	1	1.02 ± 1.32	0.12	0.94	n	y	4.64	1436	33	1042	11	y
55852	+4°2466	y	4,5	0.96 ± 1.83	0.43	0.81	n	n	3.38	6191	41	4592	2	y
	+3°2688	y	5	-	-	-	-	-	0.45	986	7	-	2	n?
	CS 22942-019	y	10	-	-	-	-	-	3.42	3274	15	2800	10	y
	CS 22948-027	y	10	-	-	-	-	-	1.87	2560	13	505	10	y

References to the Table:
(1) Začs et al. (2000) (2) This work (3) Junqueira & Pereira (2001) (4) Burris et al. (2000) (5) Luck & Bond (1991) (6) Udry et al. (1998) (10) Preston & Sneden (2001) (11) Začs et al. (2005)

4.1. Radial-velocity variations

Several stars from Table 5 (namely HIP 4347, HIP 29740, HIP 34795, HIP 55852, HIP 58956, HIP 76605 and BD +3°2688) have been monitored for many years using the CORAVEL spectrovelocimeter (Baranne et al., 1979), as part of a larger program aiming at finding the frequency of spectroscopic binaries among s-process-rich late-type giants (see Jorissen & Mayor, 1988, 1992; Jorissen et al., 1998, for details and other results from this CORAVEL monitoring). Individual radial-velocity measurements for those stars (in the CORAVEL-ELODIE system as defined in Udry et al., 1999) are given in Tables 7 or in Udry et al. (1998). For a few other stars (HIP 43042, CS 22942-019 and CS 22948-027), radial velocities were monitored using other instruments, and their results were taken from the literature (Preston & Sneden, 2001; Začs et al., 2005).

Orbits were already available for HIP 4347 and HIP 29740 (Udry et al., 1998), HIP 43042 (Začs et al., 2005), as well as for CS 22942-019 and CS 22948-027 (Preston & Sneden, 2001). A new orbital solution has been derived for BD +04°2466 (Table 6 and Fig. 8). The binary nature of those stars is therefore beyond doubt.

The radial-velocity standard deviation of HIP 58596 is larger than expected based on the uncertainty on one measurement (Fig. 9), but no satisfactory orbital solution could be found. A 3-d orbit (with an eccentricity of 0.30) is possible, but this short orbital period is not consistent with the giant nature of HIP 58596, which imposes orbital periods of at least 20 d (see Fig. 4 of Pourbaix et al., 2004). The large standard deviation exhibited by HIP 58596 is therefore very likely another example of the large intrinsic jitter often observed for metal-deficient stars, as discussed by McClure (1984) and Carney et al. (2003). Nevertheless, the proper motion analysis presented in Sect. 4.3 hints at a possible long orbital period for HIP 58596.

A 14 y radial-velocity monitoring for HIP 34795 with the northern and southern CORAVELs is not very conclusive either, mostly because there are difficulties in finding the zero-point offset between the two instruments for such large radial velocities (Udry et al., 1999). When a -1 km s^{-1} offset is applied to the northern velocities (with respect to the values listed in Table 7), a long-term trend seems to be present, albeit with some superimposed jitter (Fig. 11). The analysis of the Hipparcos astrometric

Table 4. Abundances for HD 139409, in the scale where $\log \epsilon(\mathrm{H}) = 12$, and normalized with respect to the solar abundances (Grevesse & Sauval, 1998). The standard deviations σ and the number N of lines used in the analysis are also given.

Element(X)	Z	$\log \epsilon$(X)	σ	N	[X/Fe]
C I	6	8.46	0.10	4	+0.36
Na I	11	6.02	0.05	5	+0.11
Mg I	12	7.31	0.12	6	+0.15
Si I	14	7.33	0.06	20	+0.20
Ca I	20	6.20	0.07	6	+0.26
Sc II	21	2.79	0.06	10	+0.04
Ti I	22	4.77	0.09	51	+0.17
Ti II	22	4.72	0.11	4	+0.12
V I	23	3.76	0.11	23	+0.18
Cr I	24	5.21	0.11	11	−0.04
Cr II	24	5.36	0.09	3	+0.11
Mn I	25	5.13	0.11	9	+0.16
Fe I	26	7.08	0.08	136	-
Fe II	26	7.08	0.12	14	-
Ni I	28	5.86	0.07	44	+0.03
Y I	39	2.22	0.10	2	+0.40
Y II	39	2.41	0.07	6	+0.59
Zr I	40	2.58	0.09	3	+0.40
Zr II	40	2.78	0.12	3	+0.60
Ba II	56	1.89	0.12	3	+0.18
La II	57	1.05	0.12	3	+0.30
Ce II	58	1.46	0.14	5	+0.30
Nd II	60	1.38	0.09	6	+0.30

Table 6. Orbital elements for BD +04°2466.

P (d)	4592.7±51.1
e	0.286±0.02
ω (deg)	266.8±5.3
V_γ (km s^{-1})	+39.29±0.11
K (km s^{-1})	5.67±0.12
T (JD)	2 445 076.6±73.0
$f(M)$ (M$_\odot$)	0.076
$a_1 \sin i$ (Gm)	343.18
N	41

data presented in Sect. 4.2 suggests that the star might be binary, although the evidence is not very conclusive.

Finally, there is no sign of radial-velocity variations for BD +3°2688 (Fig. 10).

4.2. Orbital motion from Hipparcos astrometric data

A tailored reprocessing of the Hipparcos *Intermediate Astrometric Data* (hereafter IAD; van Leeuwen & Evans, 1998) makes it possible to look for a possible orbital signature in the astrometric motion, following the method outlined by Pourbaix & Jorissen (2000), Pourbaix & Boffin (2003), Pourbaix (2004) and applied to barium stars by Jorissen et al. (2004b). We give here only a brief summary of the method.



Fig. 8. Radial velocities as a function of heliocentric Julian Day for BD +4°2466, superimposed on the orbital solution corresponding to the orbital elements listed in Table 6.

The basic idea is to quantify the likelihood of the fit of the Hipparcos astrometric data with an orbital model. For that purpose, Pourbaix & Arenou (2001) (see also Jancart et al., 2005) introduced several statistical indica-



Fig. 9. Radial velocities as a function of heliocentric Julian Day for HIP 58596 = HD 104340 (open circles correspond to measurements obtained with the southern CORAVEL, and filled circles with the northern one).



Fig. 10. Radial velocities as a function of heliocentric Julian Day for BD +3°2688.

tors which allow us to decide whether to keep or to discard an orbital solution. Those indicators relevant to our purpose are the following:

- The addition of 4 supplementary parameters (the four Thiele-Innes orbital constants) describing the orbital motion should result in a statistically significant decrease of the χ^2 for the fit of the N IAD with an orbital model with 9 free parameters (χ^2_T), as compared to a fit with a single-star solution with 5 free parameters (χ^2_S). This criterion is expressed by an F-test:

  $$Pr_2 = Pr[\hat{F} > F(4, N-9)], \tag{1}$$

  where

  $$\hat{F} = \frac{N-9}{4} \; \frac{\chi^2_S - \chi^2_T}{\chi^2_T}. \tag{2}$$

  Pr_2 is thus the first kind risk associated with the rejection of the null hypothesis: "*there is no orbital wobble present in the data*".
- Getting a substantial reduction of the χ^2 with the Thiele-Innes model does not necessarily imply that the four Thiele-Innes constants A, B, F, G are significantly different from 0. The first kind risk associated with the rejection of the null hypothesis "*the orbital semi-major axis is equal to zero*" may be expressed as

  $$Pr_3 = Pr[\chi^2_{ABFG} > \chi^2(4)], \tag{3}$$

  where

  $$\chi^2_{ABFG} = \boldsymbol{X}^t \boldsymbol{V}^{-1} \boldsymbol{X}, \tag{4}$$

  and $\boldsymbol{X}$ is the vector of components A, B, F, G and $\boldsymbol{V}$ is its covariance matrix.[2]
- For the orbital solution to be a significant one, its parameters should not be strongly correlated with the other astrometric parameters (e.g., the proper motion). In other words, the covariance matrix of the astrometric solution should be dominated by its diagonal terms, as measured by the *efficiency* ϵ of the matrix being close to 1 (Eichhorn, 1989). The efficiency is simply expressed by

  $$\epsilon = \sqrt[m]{\frac{\Pi_{k=1}^m \lambda_k}{\Pi_{k=1}^m \boldsymbol{V}_{kk}}}, \tag{5}$$

  where λ_k and $\boldsymbol{V}_{kk}$ are respectively the eigenvalues and the diagonal terms of the covariance matrix $\boldsymbol{V}$.

With the above notations, the requirements for a star to qualify as a binary is then

$$\alpha \equiv (Pr_2 + Pr_3)/\epsilon \leq 0.02, \tag{6}$$

where the threshold value of 0.02 has been chosen to minimize false detections (Jorissen et al., 2004b).



Fig. 11. Radial velocities as a function of heliocentric Julian Day for HIP 34795 = HD 55496. The symbols are as in Fig. 9. With respect to the data listed in Table 7, an offset of -1 km s^{-1} has been applied to the northern measurements to make them consistent with the southern ones.

---

[2] Since it may be shown that $\chi^2_S - \chi^2_T = \chi^2_{ABFG}$, the Pr_2 and Pr_3 tests are in fact equivalent provided that $\chi^2_T \sim N-9$. Thus, if Pr_2 and Pr_3 are significantly different, it means either that the Thiele-Innes orbital model does not fit the data very well ($\chi^2_T >> N-9$), or that it fits much better than could be expected ($\chi^2_T << N-9$). We are indebted to L. Lindegren for this clarification (see also Jancart et al., 2005).

Hipparcos data are, however, seldom precise enough to derive the orbital elements from scratch. Therefore, when a spectroscopic orbit is available beforehand, it is advantageous to import e, P, T from the spectroscopic orbit and to derive the remaining astrometric elements (as done by Pourbaix & Jorissen, 2000; Pourbaix & Boffin, 2003). If a spectroscopic orbit is not available, trial (e, P, T) triplets scanning a regular grid (with $10 \leq P(\mathrm{d}) \leq 5000$ imposed by the Hipparcos scanning law and the mission duration) may be used. The quality factor α is then computed for each trial (e, P, T) triplet, and if there exist triplets yielding $\alpha < 0.02$, the star is flagged as a binary. To test its success rate, this method has been applied by Jorissen et al. (2004b) on a sample of barium stars. These authors show that, when $\varpi > 5$ mas and $100 < P(\mathrm{d}) < 4000$, the (astrometric) binary detection rate is close to 100%, *i.e.*, the astrometric method recovers all known spectroscopic binaries (see also Jancart et al., 2005). When the orbit is not known beforehand, the method makes it even possible to find a good estimate for the orbital period, provided, however, that the true period is not an integer fraction, or a multiple, of one year. Here the method is applied to the sample of metal-deficient barium stars listed in Table 5.

The method flags as definite binaries the stars HIP 29740, 34795, 43042, 97874 and 107478 (Figs. 12 and 13). In two cases (HIP 29740 and 43042), the IAD method thus confirms the conclusion from the radial-velocity monitoring, but yields as well three new binaries (HIP 34795, 97874 and 107478). Two spectroscopic binaries (HIP 4347 and HIP 55852) are not detected by the IAD method because of their small parallax or long orbital period. The non-binary nature of HIP 58596, already suspected from the radial-velocity data, is confirmed by the analysis of the IAD (Fig. 13).

4.3. Orbital motion from a comparison of Hipparcos and Tycho-2 proper motions

Kaplan & Makarov (2003) suggested that the comparison of Hipparcos and Tycho-2 (Høg et al., 2000b) proper motions offers a way to detect binaries with long periods (typically from 2000 to 4000 d). The Hipparcos proper motion, being based on observations spanning only 3 y, may be altered by the orbital motion, especially for systems with periods in the range of 2000 to 4000 d whose orbital motion was not recognized by Hipparcos. On the other hand, this effect should average out in the Tycho-2 proper motion, which is derived from observations covering a much longer time span. This method, already used by Makarov (2004), Pourbaix (2004) and Jancart et al. (2006), works best when applied to stars with parallaxes in excess of about 5 mas.

The method evaluates the quantity

$$\chi^2_{\rm obs} = (\boldsymbol{\mu}_{\rm HIP} - \boldsymbol{\mu}_{\rm Tyc})^t\ \mathbf{W}^{-1}\ (\boldsymbol{\mu}_{\rm HIP} - \boldsymbol{\mu}_{\rm Tyc}), \qquad (7)$$

where $\boldsymbol{\mu}_{\rm HIP}$ and $\boldsymbol{\mu}_{\rm Tyc}$ are the vectors of α and δ components of the Hipparcos and Tycho-2 proper motions, respectively, and $\mathbf{W}$ is the associated 2×2 variance-covariance matrix. The covariance between $\mu_{\alpha,\rm HIP}$ and $\mu_{\delta,\rm HIP}$, as provided by field H28 of the Hipparcos catalogue (ESA, 1997) and the correlation between Tycho-2 and Hipparcos proper motions, as encapsulated in the quantity R of Table 1 of Høg et al. (2000a), have both been been considered (see Jancart et al. 2006 for details).

Since the above quantity follows a χ^2 probability distribution function with 2 degrees of freedom, it is then possible to compute the probability $\mathrm{Prob}(\chi^2 > \chi^2_{\rm obs})$, giving the first kind risk of rejecting the null hypothesis $\boldsymbol{\mu}_{\rm Tycho} = \boldsymbol{\mu}_{\rm HIP}$ while it is actually true. This probability is listed in Table 5, along with $\chi^2_{\rm obs}$, and the star is flagged as binary if $\mathrm{Prob} < 0.1$.

Only HIP 29740, HIP 76605 and HIP 97874 satisfy the test at the 10% threshold. Note, however, that all the other stars have parallaxes smaller than 5 mas, which make the test less efficient.

5. Summary of the binary criteria and discussion

The situation may be summarized as follows (see also last column of Table 5):

- Definite binaries with known orbits: HIP 4347, HIP 29740 (passes all three binarity tests), HIP 43042, HIP 55852, CS 22942-019, CS 22948-027;
- Suspected binaries from astrometric data (either IAD or proper motions; no or inconclusive radial-velocity data): HIP 34795, HIP 76605, HIP 97874 (both astrometric tests yield positive results), HIP 107478;
- Data inconclusive (mainly because of too small a parallax): HIP 11595, HIP 25161, HIP 69834;
- Non-binary stars (mainly from radial-velocity data, not contradicted by astrometry): HIP 58596, BD +3°2688.

The latter two non-binary stars are in fact good candidate thermally-pulsing AGB stars, as revealed by their location in the HR diagram (Table 2 and right panel of Fig. 1). Hence, they must not be binaries.

Finally, we come to the central question of this paper: Why do the metal-deficient barium stars, despite being binaries and occupying the same location of the HR diagram as YSyS, do not exhibit symbiotic activity? Three possible answers have been suggested in this paper:
(i) Some among the stars listed in Table 2 and displayed in Fig. 1 are in fact not barium stars (especially HIP 80843 = HD 148897).
(ii) Among those which are barium stars, some seem to lie on the TP-AGB, and thus need not be binaries (HIP 58596 = HD 104340, BD +3°2688). They therefore cannot exhibit symbiotic activity.
(iii) Finally, there remain a few genuine metal-deficient barium stars in the sample. Why are they not symbiotic stars? It seems that the answer to that question lies in the different period distributions for YSyS and metal-deficient barium stars: YSyS have shorter orbital period than metal-deficient barium stars, as seen in Fig. 14.



Fig. 12. The α statistics (Eq. 6) as a function of the trial orbital period (assuming $e = 0$) for metal-deficient barium stars. The period from the spectroscopic orbit (represented by an arrow) of HIP 29740 (upper left panel) indeed lies within the range of minimum α values. By comparison, the suspected binaries HIP 97874 and HIP 107478 are likely to have periods $P \sim 800$ d and $\sim 250 - 450$ d, respectively. For comparison, stars non-flagged as binaries have no (e, P, T) grid points with $\alpha < 0.02$ (see Fig. 13). The vertical dashed lines represent multiple, or integer fractions, of 1 y. At those periods, there is a strong correlation between the parallactic and orbital signals, which degrades the α statistics and makes binaries difficult to find at those 1-y alias periods.

This argument seems to apply especially to HIP 29740 (=HD 43389), which has been assigned a very bright absolute visual magnitude of −3.5 by the maximum likelihood method of Mennessier et al. (1997). It is therefore expected to have a rather strong mass loss rate, and be a good candidate YSyS. However, with its orbital period of 1689 d, it lies at the long-period edge of the period distribution of YSyS (Fig. 14). The same difference seems to exist between the period distributions of red symbiotics and binary S stars (Van Eck & Jorissen, 2002, and Fig. 14).

However, a firm conclusion on this issue should await the determination of the orbital periods for the metal-deficient barium stars flagged as binaries by the IAD



Fig. 13. Same as Fig. 12 for HIP 34795 (left panel, flagged as binary) and HIP 58596 (right panel, non-binary).

method (especially HIP 11595, HIP 25161, HIP 69834, HIP 97874, HIP 107478), so as to make the comparison between the orbital period distributions of metal-deficient barium stars and YSyS more meaningful.

Acknowledgements. This work was performed in the framework of the NATO Collaborative Linkage Grant SA (PST.CLG.979128)6774/FP. We thank D. Pourbaix for the processing of the Hipparcos data of the metal-deficient stars. F. Carrier and X. Bonfils are thanked for obtaining the Arcturus and HD 139409 spectra. LZ thanks I. Platais for valuable discussions and support. The *Fonds National Suisse de la Recherche Scientifique* has funded the operations of the CORAVEL spectrometer and of the Swiss 1-m telescope installed at the Haute-Provence Observatory.

References

Allen D. A. 1982, in M. Friedjung, R. Viotti (eds.), The Nature of Symbiotic Stars (IAU Coll. 70), Reidel, Dordrecht, 27

Allende Prieto C., García López R. J., Lambert D. L., Gustafsson B. 1999, ApJ, 527, 879

Baranne A., Mayor M., Poncet J. L. 1979, Vistas in Astronomy, 23, 279

Barbuy B., Jorissen A., Rossi S. C. F., Arnould M. 1992, A&A, 262, 216

Bessell M. S., Castelli F., Plez B. 1998, A&A, 333, 231

Boyle R. J., McClure R. D. 1975, PASP, 87, 17

Burris D. L., Pilachowski C. A., Armandroff T. E., Sneden C., Cowan J. J., Roe H. 2000, ApJ, 544, 302

Carney B. W., Latham D. W., Stefanik R. P., Laird J. B., Morse J. A. 2003, AJ, 125, 293

Clayton D. D. 1988, MNRAS, 234, 1

Corradi R., Schwarz H. E. 1997, in Physical Processes in Symbiotic Binaries and Related Systems, Copernicus Foundation for Polish Astronomy, Warsaw, 147

Corradi R. L. M., Brandi E., Ferrer O. E., Schwarz H. E. 1999, A&A, 343, 841

Drake S. A., Simon T., Linsky J. L. 1987, AJ, 93, 163

Eichhorn H. 1989, Bull. Astron. Inst. Czechosl., 40(6), 394

ESA 1997, The Hipparcos and Tycho Catalogues, ESA

Galazutdinov G. 1992, Special Astrophysical Obs. Prepr., 92, 52

Grauer A. D., Bond H. E. 1981, PASP, 93, 630

Grevesse N., Sauval A. J. 1998, Space Sci. Rev., 85, 161

Griffin R. F., Keenan P. C. 1992, The Observatory, 112, 168

Gustafsson B., Bell R. A., Eriksson K., Nordlund A. 1975, A&A, 42, 407

Hinkle K., Wallace L., Valenti J., Harmer D. 2000, Visible and Near Infrared Atlas of the Arcturus Spectrum 3727-9300 Å, Astron. Soc. Pacific, San Francisco

Høg E., Fabricius C., Makarov V. V., Bastian U., Schwekendiek P., Wicenec A., Urban S., Corbin T., Wycoff G. 2000a, A&A, 357, 367

Høg E., Fabricius C., Makarov V. V., Urban S., Corbin T., Wycoff G., Bastian U., Schwekendiek P., Wicenec A. 2000b, A&A, 355, L27

Jancart S., Jorissen A., Babusiaux C., Pourbaix D. 2005, A&A, in press

Jancart S., Jorissen A., Babusiaux C., Pourbaix D. 2006, A&A, in preparation

Jeffries R. D., Stevens I. R. 1996, MNRAS, 279, 180

Jorissen A. 2003a, in R. L. M. Corradi, J. Mikołajewska, T. J. Mahoney (eds.), Symbiotic stars probing stellar evolution, Astron. Soc. Pacific Conf. Ser. Vol. 303, San



Fig. 14. Comparison of the period distributions for samples of binary systems with different kinds of red giant primaries: S stars (Jorissen et al., 1998), red SyS (excluding symbiotic novae and symbiotic Miras; Mürset & Schmid, 1999), M giants (Jorissen et al., 2004a) and K giants (Mermilliod, 1996). The lower two panels present the orbital-period distribution for barium stars (Jorissen et al., 1998) and yellow SyS (Mürset & Schmid, 1999). In the S star panel, the arrow marked H_α indicates the period range where binary S stars exhibit H_α emission as a signature of weak symbiotic activity. The shaded area marks stars with only a lower limit available on their orbital period. The numbers in the upper right corner of each panel correspond to the sample size

Francisco, 25
Jorissen A. 2003b, in H. Habing, H. Olofsson (eds.), Asymptotic Giant Branch Stars, Springer Verlag, New York, p. 461
Jorissen A., Famaey B., Dedecker M., Pourbaix D., Mayor M., Udry S. 2004a, Rev. Mex. Astron. Astrof. Conf. Ser., 71–72
Jorissen A., Jancart S., Pourbaix D. 2004b, in R. W. Hilditch, H. Hensberge, K. Pavlovski (eds.), Spectroscopically and Spatially Resolving the Components of Close Binary Stars, San Francisco: Astron. Soc. Pacific Conf. Ser. 318, 141
Jorissen A., Mayor M. 1988, A&A, 198, 187
Jorissen A., Mayor M. 1992, A&A, 260, 115
Jorissen A., Schmitt J. H. M. M., Carquillat J. M., Ginestet N., Bickert K. F. 1996, A&A, 306, 467
Jorissen A., Van Eck S., Mayor M., Udry S. 1998, A&A, 332, 877
Junqueira S., Pereira C. B. 2001, AJ, 122, 360
Kaplan G. H., Makarov V. V. 2003, Astronomische Nachrichten, 324, 419
Keenan P. C., McNeil R. C. 1989, ApJS, 71, 245
Kyröläinen J., Tuominen I., Vilhu O., Virtanen H. 1986, A&AS, 65, 11
Lattanzio J. C. 1991, ApJS, 76, 215
Lü P. K. 1991, AJ, 101, 2229
Lü P. K., Dawson D. W., Upgren A. R., Weis E. W. 1983, ApJS, 52, 169
Luck R. E. 1991, ApJS, 75, 579
Luck R. E., Bond H. E. 1991, ApJS, 77, 515
MacConnell D. J., Frye R. L., Upgren A. R. 1972, AJ, 77, 384
Makarov V. V. 2004, ApJ, 600, L71
Mayor M., Pepe F., Queloz D., Bouchy F., Rupprecht G., Lo Curto G., Avila G., Benz W., Bertaux J.-L., Bonfils X., dall T., Dekker H., Delabre B., Eckert W., Fleury M., Gilliotte A., Gojak D., Guzman J. C., Kohler D., Lizon J.-L., Longinotti A., Lovis C., Megevand D., Pasquini L., Reyes J., Sivan J.-P., Sosnowska D., Soto R., Udry S., van Kesteren A., Weber L., Weilenmann U. 2003, The Messenger, 114, 20
McClure R. D. 1984, ApJ, 280, L31
McWilliam A. 1997, Ann. Rev. Astron. Astrophys., 35, 503
Mennessier M. O., Luri X., Figueras F., Gomez A. E., Grenier S., Torra J., North P. 1997, A&A, 326, 722
Mermilliod J.-C. 1996, in E. F. Milone, J.-C. Mermilliod (eds.), The Origins, Evolution and Destinies of Binary Stars in Clusters (ASP Conf. Ser. 90), Astron. Soc. Pacific, San Francisco, 95
Munari U., Buson L. M. 1992, A&A, 255, 158
Munari U., Yudin B. F., Taranova O. G., Massone G., Marang F., Roberts G., Winkler H., Whitelock P. A. 1992, A&AS, 93, 383
Mürset U., Nussbaumer H., Schmid H. M., Vogel M. 1991, A&A, 248, 458
Mürset U., Schmid H. M. 1999, A&AS, 137, 473
Musaev F. A., Galazutdinov G. A., Sergeev A. V., Karpov N. V., Podyachaev Y. V. 1999, Kinematika I Fizika Nebesnyh Tel, 15, 282
Pereira C. B., Porto de Mello G. F. 1997, AJ, 114, 2128
Pereira C. B., Smith V. V., Cunha K. 1998, AJ, 116, 1977
Pinsonneault M. H., Sneden C., Smith V. V. 1984, PASP, 96, 239
Piskunov N. E., Kupka F., Ryabchikova T. A., Weiss W. W., Jeffery C. S. 1995, A&AS, 112, 525
Pourbaix D. 2004, in R. W. Hilditch, H. Hensberge, K. Pavlovski (eds.), Spectroscopically and Spatially Resolving the Components of Close Binary Stars, San Francisco: Astron. Soc. Pacific Conf. Ser. 318, p. 132
Pourbaix D., Arenou F. 2001, A&A, 372, 935
Pourbaix D., Boffin H. M. J. 2003, A&A, 398, 1163
Pourbaix D., Jorissen A. 2000, A&AS, 145, 161
Pourbaix D., Tokovinin A. A., Batten A. H., Fekel F. C., Hartkopf W. I., Levato H., Morrell N. I., Torres G., Udry S. 2004, A&A, 424, 727
Preston G. W., Sneden C. 2001, AJ, 122, 1545
Schaller G., Schaerer D., Meynet G., Maeder A. 1992, A&AS, 96, 269

Schmid H. M., Nussbaumer H. 1993, A&A, 268, 159

Schwarz H. E. 1991, A&A, 243, 469

Šleivyte J., Bartkevičius A. 1990, Vilnius Astronomijos Observatorijos Biuletenis, 85, 3

Smith V. V., Cunha K., Jorissen A., Boffin H. M. J. 1996, A&A, 315, 179

Smith V. V., Cunha K., Jorissen A., Boffin H. M. J. 1997, A&A, 324, 97

Theuns T., Boffin H. M. J., Jorissen A. 1996, MNRAS, 280, 1264

Tsymbal V. 1996, in S. J. Adelman, F. Kupka, W. W. Weiss (eds.), Model Atmospheres and Spectrum Synthesis, ASP Conf. Ser. 108, Astron. Soc. Pacific, San Francisco, 198

Udry S., Jorissen A., Mayor M., Van Eck S. 1998, A&AS, 131, 25

Udry S., Mayor M., Queloz D. 1999, in J. B. Hearnshaw, C. D. Scarfe (eds.), Precise Stellar Radial Velocities, Astron. Soc. Pacific Conf. Ser., Vol.170, San Francisco, 367

Van Eck S., Jorissen A. 2002, A&A, 396, 599

van Leeuwen F., Evans D. W. 1998, A&AS, 130, 157

Vilhu O., Tuominen I., Kyrolainen J., Holsti N. 1977, Obs. and Astrophys. Lab. Univ. Helsinki Rep. No 3

Yamashita Y., Norimoto Y. 1981, Annals of the Tokyo Astronomical Observatory, 18, 125

Začs L., Schmidt M. R., Musaev, F. A. Galazutdinov G. A., Sperauskas J. 2005, A&A, in press

Začs L., Schmidt M. R., Schuster W. J. 2000, A&A, 358, 1022

Table 7. Individual CORAVEL radial velocities and associated errors for HIP 34795, HIP 58596, HIP 76605, BD +3°2688 and BD +04°2466. The last column indicates which one of the two CORAVEL spectrovelocimeters has been used (NO = CORAVEL north at the *Observatoire de Haute Provence*; SO = CORAVEL south on the Danish 1.54-m telescope at ESO)

HIP 34795 = HD 55496

DDMMYY	HJD	R.V. (km s^{-1})	ϵ (km s^{-1})	
050183	45340.695	315.30	0.33	SO
151283	45684.797	314.79	0.33	SO
120684	45864.434	315.71	0.46	SO
310185	46097.658	314.78	0.34	SO
271185	46397.826	314.95	0.30	SO
111186	46746.858	314.56	0.31	SO
101286	46775.595	316.07	0.56	NO
150187	46811.682	314.98	0.31	SO
160288	47208.394	316.01	0.49	NO
231088	47458.704	315.83	0.70	NO
151288	47511.857	315.11	0.33	SO
090289	47567.393	315.39	0.56	NO
241189	47855.642	315.44	0.57	NO
070390	47958.339	317.36	0.66	NO
050490	47987.497	314.78	0.31	SO
061190	48202.697	315.63	0.45	NO
031290	48229.645	316.10	0.56	NO
180391	48334.380	314.95	0.62	NO
110393	49058.361	315.00	0.67	NO
130393	49060.338	314.32	0.80	NO
180194	49371.471	315.37	0.58	NO
301295	50082.752	313.99	0.34	SO
231196	50411.600	316.66	0.56	NO
120197	50461.480	317.72	0.63	NO

HIP 58596 = HD 104340

DDMMYY	HJD	R.V. (km s^{-1})	ϵ (km s^{-1})	
141289	47875.832	265.29	0.45	SO
140392	48696.805	264.18	0.30	SO
120393	49059.755	266.40	0.29	SO
190393	49066.707	264.89	0.33	SO
050493	49083.708	265.41	0.28	SO
200493	49098.453	263.93	0.46	NO
080394	49420.525	263.34	0.47	NO
030494	49446.435	266.38	0.41	NO
140594	49487.360	265.37	0.46	NO
260195	49744.638	264.59	0.56	NO
300395	49807.475	263.16	0.52	NO
210795	49920.516	264.34	0.30	SO
191295	50071.721	265.60	0.37	NO
150296	50129.556	263.70	0.43	NO
130197	50462.684	265.61	0.45	NO

HIP 76605 = HD 139409

DDMMYY	HJD	R.V. (km s^{-1})	ϵ (km s^{-1})	
090790	48082.629	62.43	0.30	SO
260794	49560.513	63.36	0.27	SO

Table 7. Continued.

HIP 55852 = BD +04°2466

DDMMYY	HJD	R.V. (km s^{-1})	ϵ (km s^{-1})	
070282	45008.814	38.11	0.69	SO
130684	45865.491	44.73	0.74	SO
030285	46100.832	45.14	0.68	SO
290385	46154.675	44.78	0.58	SO
200186	46451.831	43.54	0.53	SO
140286	46476.783	43.32	0.49	SO
291186	46764.706	42.05	0.96	NO
090187	46805.864	41.36	0.49	SO
080287	46835.808	41.99	0.41	SO
280188	47189.812	40.50	0.43	SO
040388	47225.726	39.83	0.41	SO
250488	47277.657	40.17	0.48	SO
051288	47501.707	37.84	0.69	NO
060389	47592.565	38.14	0.64	NO
220589	47669.526	37.64	0.49	SO
241289	47885.857	37.48	0.45	SO
050290	47928.800	36.83	0.48	SO
160290	47939.743	37.05	0.43	SO
260290	47949.595	37.00	0.69	NO
040490	47986.631	37.01	0.42	SO
110490	47993.594	35.99	0.49	SO
050590	48017.647	36.35	0.61	SO
020690	48045.501	36.82	0.46	SO
260291	48314.487	34.56	0.73	NO
260391	48342.724	34.92	0.47	SO
150392	48697.747	34.26	0.42	SO
120393	49059.726	33.77	0.39	SO
070493	49085.670	33.15	0.40	SO
080493	49086.448	32.60	0.70	NO
100294	49394.609	34.54	0.77	NO
020394	49414.713	35.84	0.43	SO
070394	49419.515	34.92	0.52	NO
080494	49451.424	37.51	0.88	NO
140594	49487.351	36.78	0.64	NO
310195	49749.592	39.24	0.74	NO
150395	49792.681	41.24	0.44	SO
020495	49810.460	40.55	0.44	NO
190795	49918.506	41.94	0.53	SO
191295	50071.715	43.32	0.60	NO
130197	50462.674	44.33	0.90	NO
200199	51199.634	41.42	0.80	NO

BD +03°2688

DDMMYY	HJD	R.V. (km s^{-1})	ϵ (km s^{-1})	
080493	49086.481	33.16	0.43	NO
050394	49417.506	32.35	0.47	NO
030494	49446.510	32.33	0.44	NO
140594	49487.388	33.00	0.47	NO
030295	49752.658	33.52	0.52	NO
300395	49807.506	32.64	0.71	NO
201295	50072.729	33.07	0.46	NO